WHEN TO REFER: Guidelines for a Speech Pathology Referral

Age	Language	Speech	Social/Emotional	Feeding
0 - 18 months	 Child is a quiet baby and does not babble much. Child is unresponsive to familiar people/words. 	Hearing difficulties.Child is prone to colds or ear infections.	 Lacks eye contact. Does not show social enjoyment. Lacks interest in playing and social interactions. 	 Difficulties with bottle feeding and or breast feeding. Child has swallowing difficulties. Failure to thrive. Difficulty transitioning to solids.
2 - 3 years	 Child has less than 50 words in their vocabulary. Child is not putting 2 words together. Child has difficulty following simple instructions. 	 Child has no clear words, and is difficult to understand when speaking. Child's speech is dysfluent, or 'bumpy'. 	 Lacks eye contact. Plays with toys inappropriately. Such as banging/throwing doll, instead of cuddling doll. Lacks interest in pretend play, or playing with others. Child has difficulties showing different feelings. 	Child is a fussy eater, and reluctant to try need food or textures.
3 – 5 years	 Child is not using simple sentences. Child has difficulty answering simple questions. Difficulty expressing needs/wants/telling a story. Difficulty following 2 step instructions. 	 Most of what is said cannot be understood by other person. Child's speech is dysfluent, or 'bumpy'. Child has a 'harsh' or 'hoarse' sounding voice. 	 Lacks eye contact. Does not play cooperatively with others. Difficulties in noticing and understanding feelings of other people. 	BRIGH